



Somatostatin Receptor Type 2 (SSTR2) Antibody


Catalogue No.: abx317864




Western blot analysis of Mouse spleen tissue using SSTR2 Antibody (2.7 µg/ml) and Goat Polyclonal Anti-Rabbit IgG (1/50000 dilution). Calculated MW: 42 kDa, 41 kDa. Observed MW: 42 kDa.



IHC-P analysis of Human kidney tissue using SSTR2 Antibody (1/100 dilution).



IF analysis of HepG2 cells using SSTR2 Antibody (1/100 dilution) and AF488-conjugated Goat Anti-Rabbit IgG.



IF analysis of PC-3 cells using SSTR2 Antibody (1/100 dilution) and AF488-conjugated Goat Anti-Rabbit IgG.

SSTR2 Antibody is a Rabbit Polyclonal against SSTR2.

Target: Somatostatin Receptor Type 2 (SSTR2)

Clonality: Polyclonal

Reactivity: Human, Mouse

Datasheet

Version: 4.0.0
Revision date: 25 Sep 2025



Tested Applications: ELISA, WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 1/1000 - 1/5000, IHC: 1/20 - 1/200, IF/ICC: 1/50 - 1/200. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: Recombinant human Somatostatin receptor type 2 protein (304-369 AA).

Isotype: IgG

Form: Liquid

Purity: > 95%

Purification: Purified by Protein G.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: P30874 ([UniProt](#), [ExPASy](#))

Gene Symbol: SSTR2

GeneID: [6752](#)

OMIM: [182452](#)

NCBI Accession: NP_001041.1

KEGG: hsa:6752

String: [9606.ENSP00000350198](#)

Molecular Weight: Calculated MW: 42 kDa, 41 kDa
Observed MW: 42 kDa

Buffer: 0.01 M PBS, pH 7.4, containing 0.03% Proclin-300 and 50% Glycerol.

Concentration: 4 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.