


Receptor Interacting Serine Threonine Kinase 2 (RIPK2) Antibody

Catalogue No.: abx001942



Western blot analysis of various lysates using RIPK2 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 90s.

RIPK2 Antibody is a Rabbit Polyclonal antibody against RIPK2. This gene encodes a member of the receptor-interacting protein (RIP) family of serine/threonine protein kinases. The encoded protein contains a C-terminal caspase activation and recruitment domain (CARD), and is a component of signaling complexes in both the innate and adaptive immune pathways. It is a potent activator of NF-κB and inducer of apoptosis in response to various stimuli.

Target:	Receptor Interacting Serine Threonine Kinase 2 (RIPK2)
Clonality:	Polyclonal
Reactivity:	Human, Mouse, Rat
Tested Applications:	ELISA, WB
Host:	Rabbit
Recommended dilutions:	ELISA: 1 µg/ml, WB: 1/500 - 1/1000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant protein corresponding to RIPK2. The exact sequence is proprietary.
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	O43353 (UniProt , ExPASy)
Gene Symbol:	RIPK2

Datasheet

Version: 5.0.0

Revision date: 01 Oct 2025



GeneID: [8767](#)

NCBI Accession: NP_003812.1

KEGG: hsa:8767

String: [9606.ENSP00000220751](#)

Molecular Weight: Calculated MW: 61 kDa
Observed MW: 61 kDa

Buffer: PBS, pH 7.3, containing 0.05% Proclin-300, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only