Datasheet

Version: 3.0.0 Revision date: 04 Jun 2025



B-Cell Linker Protein (BLNK) Antibody

Catalogue No.:abx025447



This gene encodes a cytoplasmic linker or adaptor protein that plays a critical role in B cell development. This protein bridges B cell receptor-associated kinase activation with downstream signaling pathways, thereby affecting various biological functions. The phosphorylation of five tyrosine residues is necessary for this protein to nucleate distinct signaling effectors following B cell receptor activation. Mutations in this gene cause hypoglobulinemia and absent B cells, a disease in which the pro to pre-B-cell transition is developmentally blocked. Deficiency in this protein has also been shown in some cases of pre-B acute lymphoblastic leukemia. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. This antibody is supplied as crude ascites.

Target: B-Cell Linker Protein (BLNK)

Clonality: Monoclonal

Reactivity: Human

Tested Applications: ELISA, WB

Host: Mouse

Recommended dilutions: WB: 1/500 - 1/8000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: KLH-conjugated synthetic peptide between 150-178 amino acids from human BLNK.

Isotype: IgM

Form: Liquid

Purification: Unpurified crude ascites.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q8WV28 (<u>UniProt</u>, <u>ExPASy</u>)

Website: www.abbexa.com · Email: info@abbexa.com

Datasheet

Version: 3.0.0 Revision date: 04 Jun 2025



KEGG: hsa:29760

String: <u>9606.ENSP00000224337</u>

Molecular Weight: Calculated MW: 50.5 kDa

Buffer: Ascites containing 0.09% sodium azide.

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY, NOT FOR USE IN DIAGNOSTIC.

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.