



Apoptosis Regulator Bcl-2 (BCL2) Antibody


Catalogue No.: abx230840



WB analysis of U-937 cells, using BCL2 antibody (1/6000 dilution).



IHC-P analysis of human tonsillitis tissue, using BCL2 Antibody (1/100 dilution).



IP analysis of HL-60 cell lysate (4000 µg), using BCL2 antibody (4 µg, detection: 1/1000 dilution).

BCL2 Antibody is a Mouse Monoclonal against BCL2.

Target: Apoptosis Regulator Bcl-2 (BCL2)

Clonality: Monoclonal

Reactivity: Human

Tested Applications: ELISA, WB, IHC, IP

Host: Mouse

Recommended dilutions: WB: 1/1000 - 1/10000, IHC: 1/100 - 1/2000, IP: 1/500 - 1/5000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Datasheet

Version: 2.0.0
Revision date: 07 Jul 2025



Immunogen:	B-cell CLL/lymphoma 2
Isotype:	IgG _{2a}
Form:	Liquid
Purity:	≥ 95% (SDS-PAGE)
Purification:	Purified by Protein A and Protein G affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	P10415 (UniProt , ExPASy)
Gene Symbol:	BCL2
GeneID:	596
OMIM:	151430
HGNC:	990
KEGG:	hsa:596
Ensembl:	ENSG00000171791
String:	9606.ENSP00000381185
Molecular Weight:	Observed MW: 26 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.