


Glycerol-3-Phosphate Acyltransferase, Mitochondrial (GPAM) Antibody

Catalogue No.: abx025609



Glycerol-3-phosphate acyltransferase (GPAT; EC 2.3.1.15), which catalyzes the initial and committing step in glycerolipid biosynthesis, is predicted to play a pivotal role in the regulation of cellular triacylglycerol and phospholipid levels. Two mammalian forms of GPAT have been identified on the basis of localization to either the endoplasmic reticulum or mitochondria.[supplied by OMIM].

Target: Glycerol-3-Phosphate Acyltransferase, Mitochondrial (GPAM)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: ELISA, WB, FCM

Host: Rabbit

Recommended dilutions: WB: 1/1000, FCM: 1/10 - 1/50. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: KLH-conjugated synthetic peptide between 426-455 amino acids from the Central region of human GPAM.

Isotype: IgG

Datasheet

Version: 2.0.0

Revision date: 30 Jun 2025



| | |
|----------------------------|--|
| Form: | Liquid |
| Purification: | Purified through a protein A column, followed by peptide affinity purification. |
| Storage: | Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles. |
| UniProt Primary AC: | Q9HCL2 (UniProt , ExPASy) |
| Gene Symbol: | GPAM |
| String: | 9606.ENSP00000265276 |
| Molecular Weight: | Calculated MW: 93.8 kDa |
| Buffer: | PBS containing 0.09% sodium azide. |
| Note: | THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION. |

For Reference Only