



Triokinase/FMN Cyclase (DAK) Antibody

Catalogue No.: abx232233



WB analysis of HeLa cells, using DAK antibody (1/3000 dilution).



IP analysis of HeLa cell lysate (2150 µg), using DAK antibody (3 µg, detection: 1/500 dilution).

DAK Antibody is a Rabbit Polyclonal against DAK.

Target: Triokinase/FMN Cyclase (DAK)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB, IP

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000, IP: 1/200 - 1/2000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: dihydroxyacetone kinase 2 homolog(S. cerevisiae)

Isotype: IgG

Form: Liquid

Purity: ≥ 95% (SDS-PAGE)

Datasheet

Version: 2.0.0

Revision date: 10 Sep 2025



Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	Q3LXA3 (UniProt , ExPASy)
Gene Symbol:	TKFC
GeneID:	26007
OMIM:	615844
HGNC:	24552
KEGG:	hsa:26007
Ensembl:	ENSG00000149476
String:	9606.ENSP00000378360
Molecular Weight:	Observed MW: 59 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.