



Endothelial Monocyte-Activating Polypeptide 2 (EMAP II) Antibody

Catalogue No.: abx232750



WB analysis of SGC-7901 cells, using SCYE1 antibody (1/500 dilution).



IHC-P analysis of human cervical cancer tissue, using SCYE1 antibody (1/100 dilution).

EMAP II Antibody is a Rabbit Polyclonal against EMAP II.

Target: Endothelial Monocyte-Activating Polypeptide 2 (EMAP II)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000, IHC: 1/20 - 1/200, IF/ICC: 1/20 - 1/200. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: small inducible cytokine subfamily E, member 1 (endothelial monocyte-activating)

Isotype: IgG

Form: Liquid

Purity: ≥ 95% (SDS-PAGE)

Datasheet

Version: 3.0.0
Revision date: 17 Jun 2025



Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	Q12904 (UniProt , ExPASy)
KEGG:	hsa:9255
String:	9606.ENSP00000378191
Molecular Weight:	Observed MW: 34 kDa, 68 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only