



NAD(P)H Dehydrogenase, Quinone 1 (NQO1) Antibody


Catalogue No.: abx235830



WB analysis of K-562 cells, using NQO1 antibody (1/1000 dilution).



IHC-P analysis of human pancreas cancer tissue, using NQO1 Antibody (1/200 dilution).



IP analysis of HepG2 cell lysate (3000 µg), using NQO1 antibody (4 µg, detection: 1/300 dilution).

NQO1 Antibody is a Rabbit Polyclonal against NQO1.

Target: NAD(P)H Dehydrogenase, Quinone 1 (NQO1)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB, IHC, IF/ICC, FCM, IP

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000, IHC: 1/100 - 1/500, IF/ICC: 1/20 - 1/200, IP: 1/150 - 1/600. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Datasheet

Version: 1.0.0
Revision date: 01 Jun 2025



Immunogen:	NAD (P) H dehydrogenase, quinone 1
Isotype:	IgG
Form:	Liquid
Purity:	≥ 95% (SDS-PAGE)
Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	P15559 (UniProt , ExPASy)
Gene Symbol:	NQO1
Molecular Weight:	Observed MW: 31 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.