



26S Proteasome Regulatory Subunit 10B (PSMC6) Antibody

Catalogue No.:abx236881



WB analysis of mouse liver tissue, using PSMC6 antibody (1/1000 dilution).



IHC-P analysis of human kidney tissue, using PSMC6 Antibody (1/100 dilution).

26S Proteasome Regulatory Subunit 10B (PSMC6) Antibody is a Rabbit Polyclonal against 26S Proteasome Regulatory Subunit 10B (PSMC6). The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases which have a chaperone-like activity. Pseudogenes have been identified on chromosomes 8 and 12.

Target: 26S Proteasome Regulatory Subunit 10B (PSMC6)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB, IHC

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000, IHC: 1/50 - 1/200. Optimal dilutions/concentrations should be determined by

the end user.

Conjugation: Unconjugated

Immunogen: proteasome (prosome, macropain) 26S subunit, ATPase, 6

Datasheet

Version: 3.0.0 Revision date: 05 Oct 2025



Isotype: IgG

Form: Liquid

Purity: $\geq 95\%$ (SDS-PAGE)

Purification: Purified by immunogen affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Validity: 12 months.

UniProt Primary AC: P62333 (<u>UniProt</u>, <u>ExPASy</u>)

Gene Symbol: PSMC6

GeneID: <u>5706</u>

OMIM: <u>602708</u>

HGNC: 9553

KEGG: hsa:5706

Ensembl: ENSG00000100519

String: 9606.ENSP00000401802

Molecular Weight: Observed MW: 45 kDa

Buffer: PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.

Concentration: 2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC,

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.