



Proteasome Activator Subunit 4 (PSME4) Antibody


Catalogue No.: abx236897



WB analysis of HEK-293 cells, using PSME4 antibody (1/400 dilution).



IHC-P analysis of human liver cancer tissue, using PSME4 antibody (1/200 dilution).



IP analysis of HEK-293 cell lysate (1760 µg), using PSME4 antibody (3 µg, detection: 1/500 dilution).

PSME4 Antibody is a Rabbit Polyclonal against PSME4.

Target: Proteasome Activator Subunit 4 (PSME4)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB, IHC, IF/ICC, IP

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000, IHC: 1/50 - 1/500, IF/ICC: 1/10 - 1/100, IP: 1/500 - 1/1000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Datasheet

Version: 2.0.0
Revision date: 18 Mar 2025



Immunogen:	proteasome (prosome, macropain) activator subunit 4
Isotype:	IgG
Form:	Liquid
Purity:	≥ 95% (SDS-PAGE)
Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	Q14997 (UniProt , ExPASy)
Gene Symbol:	PSME4
GeneID:	23198
OMIM:	607705
HGNC:	20635
KEGG:	hsa:23198
Ensembl:	ENSG00000068878
String:	9606.ENSP00000384211
Molecular Weight:	Observed MW: 140 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.