



Recombination Activating Gene 2 (RAG2) Antibody


Catalogue No.: abx237089



WB analysis of A375 cells, using RAG2 antibody (1/300 dilution).



IHC-P analysis of human lymphoma tissue, using RAG2 antibody (1/50 dilution).



IP analysis of A375 cell lysate (3600 µg), using RAG2 antibody (4 µg, detection: 1/300 dilution).

RAG2 Antibody is a Rabbit Polyclonal against RAG2.

Target: Recombination Activating Gene 2 (RAG2)

Clonality: Polyclonal

Reactivity: Human, Mouse

Tested Applications: ELISA, WB, IHC, IF/ICC, FCM, IP

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000, IHC: 1/20 - 1/200, IF/ICC: 1/20 - 1/200, IP: 1/200 - 1/1000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Datasheet

Version: 2.0.0
Revision date: 29 Sep 2025



Immunogen:	recombination activating gene 2
Isotype:	IgG
Form:	Liquid
Purity:	≥ 95% (SDS-PAGE)
Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	P55895 (UniProt , ExPASy)
Gene Symbol:	RAG2
GeneID:	5897
OMIM:	179616
HGNC:	9832
KEGG:	hsa:5897
String:	9606.ENSP00000478672
Molecular Weight:	Observed MW: 57-62 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.