



S-Adenosylmethionine-Dependent Nucleotide Dehydratase RSAD2 (RSAD2) Antibody


Catalogue No.: abx237504



WB analysis of A549 cells, using RSAD2 antibody (1/500 dilution).



IHC-P analysis of human skin tissue, using RSAD2 antibody (1/50 dilution).



IF analysis of HeLa cells, using RSAD2 antibody (1/25 dilution) and Rhodamine-conjugated goat anti-rabbit IgG (red).

RSAD2 Antibody is a Rabbit Polyclonal against RSAD2.

Target: S-Adenosylmethionine-Dependent Nucleotide Dehydratase RSAD2 (RSAD2)

Clonality: Polyclonal

Reactivity: Human, Mouse

Tested Applications: ELISA, WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/1000, IHC: 1/20 - 1/200, IF/ICC: 1/10 - 1/100. Optimal dilutions/concentrations should be determined by the end user.

Datasheet

Version: 4.0.0
Revision date: 18 Mar 2025



Conjugation:	Unconjugated
Immunogen:	radical S-adenosyl methionine domain containing 2
Isotype:	IgG
Form:	Liquid
Purity:	≥ 95% (SDS-PAGE)
Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	Q8WVG1 (UniProt , ExPASy)
Gene Symbol:	RSAD2
GeneID:	91543
OMIM:	607810
HGNC:	30908
KEGG:	hsa:91543
Ensembl:	ENSG00000134321
String:	9606.ENSP00000371471
Molecular Weight:	Observed MW: 45 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.