



NAD-Dependent Protein Deacetylase Sirtuin-2 (SIRT2) Antibody

Catalogue No.: abx237879



WB analysis of mouse brain tissue, using SIRT2 antibody (1/1000 dilution).



IHC-P analysis of human lung cancer tissue, using SIRT2 Antibody (1/100 dilution).

SIRT2 Antibody is a Rabbit Polyclonal against SIRT2.

Target: NAD-Dependent Protein Deacetylase Sirtuin-2 (SIRT2)

Research Area: Signal Transduction, Developmental Science

Clonality: Polyclonal

Reactivity: Human, Rat

Tested Applications: ELISA, WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000, IHC: 1/50 - 1/200, IF/ICC: 1/10 - 1/100. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: sirtuin (silent mating type information regulation 2 homolog) 2

Isotype: IgG

Form: Liquid

Datasheet

Version: 5.0.0

Revision date: 06 Sep 2025



Purity:	≥ 95% (SDS-PAGE)
Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	Q8IXJ6 (UniProt , ExPASy)
KEGG:	hsa:22933
String:	9606.ENSP00000249396
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only