



NAD-Dependent Protein Deacetylase Sirtuin-3, Mitochondrial (SIRT3) Antibody

Catalogue No.: abx237881



WB analysis of SW620 cells, using SIRT3 antibody (1/1000 dilution).



IHC-P analysis of rat kidney tissue, using SIRT3 Antibody (1/100 dilution).

SIRT3 Antibody is a Rabbit Polyclonal against SIRT3.

Target: NAD-Dependent Protein Deacetylase Sirtuin-3, Mitochondrial (SIRT3)

Research Area: Signal Transduction

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000, IHC: 1/50 - 1/100, IF/ICC: 1/50 - 1/100. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: sirtuin (silent mating type information regulation 2 homolog) 3

Isotype: IgG

Datasheet

Version: 4.0.0

Revision date: 31 Aug 2025



Form:	Liquid
Purity:	≥ 95% (SDS-PAGE)
Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	Q9NTG7 (UniProt , ExPASy)
String:	9606.ENSP00000372191
Molecular Weight:	Observed MW: 44 kDa, 27 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only