


NAD-Dependent Protein Deacylase Sirtuin-5, Mitochondrial (SIRT5) Antibody

Catalogue No.: abx237883



WB analysis of HeLa cells, using SIRT5 antibody (1/600 dilution).

IP analysis of mouse liver tissue lysate (6000 µg), using SIRT5 antibody (4 µg, detection: 1/500 dilution).

SIRT5 Antibody is a Rabbit Polyclonal against SIRT5.

Target: NAD-Dependent Protein Deacylase Sirtuin-5, Mitochondrial (SIRT5)

Research Area: Signal Transduction

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB, IP

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/1000, IF/ICC: 1/50 - 1/500, IP: 1/500 - 1/1000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: sirtuin (silent mating type information regulation 2 homolog) 5

Isotype: IgG

Datasheet

Version: 3.0.0

Revision date: 19 Sep 2025



Form:	Liquid
Purity:	≥ 95% (SDS-PAGE)
Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	Q9NXA8 (UniProt , ExPASy)
String:	9606.ENSP00000476228
Molecular Weight:	Observed MW: 33 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only