



Suppressors of Cytokine Signaling 3 (SOCS3) Antibody


Catalogue No.: abx238100




WB analysis of K-562 cells, using SOCS3 antibody (1/500 dilution).



IHC-P analysis of human heart tissue, using SOCS3 antibody (1/50 dilution).



IF analysis of HepG2 cells, using SOCS3 antibody (1/10 dilution) and Rhodamine-conjugated goat anti-rabbit IgG (red). Blue pseudocolor = DAPI (fluorescent DNA dye).



IP analysis of K-562 cell lysate (1720 µg), using SOCS3 antibody (4 µg, detection: 1/500 dilution).

SOCS3 Antibody is a Rabbit Polyclonal against SOCS3.

Target: Suppressors of Cytokine Signaling 3 (SOCS3)**Clonality:** Polyclonal**Reactivity:** Human, Mouse, Rat

Datasheet

Version: 2.0.0
Revision date: 02 Oct 2025



Tested Applications:	ELISA, WB, IHC, IF/ICC, IP
Host:	Rabbit
Recommended dilutions:	WB: 1/200 - 1/2000, IHC: 1/20 - 1/200, IF/ICC: 1/10 - 1/100, IP: 1/200 - 1/2000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	suppressor of cytokine signaling 3
Isotype:	IgG
Form:	Liquid
Purity:	≥ 95% (SDS-PAGE)
Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	O14543 (UniProt , ExPASy)
Gene Symbol:	SOCS3
OMIM:	604176
HGNC:	19391
KEGG:	hsa:9021
String:	9606.ENSP00000330341
Molecular Weight:	Observed MW: 30 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.