



SRC Proto-Oncogene, Non-Receptor Tyrosine Kinase (SRC) Antibody


Catalogue No.: abx238216




WB analysis of Jurkat cells, using SRC antibody (1/1000 dilution).



IHC-P analysis of human breast cancer tissue, using c-SRC Antibody (1/200 dilution).



IF analysis of A431 cells, using SRC antibody (1/25 dilution) and AF488-conjugated Goat Anti-Rabbit IgG (H+L).



IP analysis of SH-SY5Y cell lysate (2900 µg), using SRC antibody (5 µg, detection: 1/1000 dilution).

SRC Antibody is a Rabbit Polyclonal against SRC.

Target: SRC Proto-Oncogene, Non-Receptor Tyrosine Kinase (SRC)**Clonality:** Polyclonal**Reactivity:** Human, Mouse, Rat, Hamster

Datasheet

Version: 2.0.0
Revision date: 25 Sep 2025



Tested Applications:	ELISA, WB, IHC, IF/ICC, FCM, IP
Host:	Rabbit
Recommended dilutions:	WB: 1/500 - 1/2000, IHC: 1/100 - 1/400, IF/ICC: 1/10 - 1/100, IP: 1/500 - 1/2000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	v-src sarcoma (Schmidt-Ruppin A-2) viral oncogene homolog (avian)
Isotype:	IgG
Form:	Liquid
Purity:	≥ 95% (SDS-PAGE)
Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	P12931 (UniProt , ExPASy)
Gene Symbol:	SRC
GeneID:	6714
OMIM:	190090
HGNC:	11283
Ensembl:	ENSG00000197122
Molecular Weight:	Observed MW: 55 kDa, 60 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.