



Aldehyde Dehydrogenase 4 Family, Member A1 (ALDH4A1) Antibody

Catalogue No.:abx002029



Western blot analysis of various lysates, using ALDH4A1 Antibody at 1/6000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 10s



Immunofluorescence analysis of A549 cells using ALDH4A1 Antibody. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1/500 dilution.

ALDH4A1 Antibody is a Rabbit Polyclonal antibody against ALDH4A1. This protein belongs to the aldehyde dehydrogenase family of proteins. This enzyme is a mitochondrial matrix NAD-dependent dehydrogenase which catalyzes the second step of the proline degradation pathway, converting pyrroline-5-carboxylate to glutamate. Deficiency of this enzyme is associated with type II hyperprolinemia, an autosomal recessive disorder characterized by accumulation of delta-1-pyrroline-5-carboxylate (P5C) and proline. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene.

Target: Aldehyde Dehydrogenase 4 Family, Member A1 (ALDH4A1)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB, IF/ICC

Host: Rabbit

Recommended dilutions: ELISA: 1 µg/ml, WB: 1/2000 - 1/7000, IF/ICC: 1/50 - 1/200. Optimal dilutions/concentrations should

be determined by the end user.

Conjugation: Unconjugated

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 354-563 of

human ALDH4A1.

Isotype: IgG

Datasheet

Version: 3.0.0 Revision date: 28 Jun 2025



Form: Liquid

Purification: Purified by affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: P30038 (<u>UniProt</u>, <u>ExPASy</u>)

Gene Symbol: ALDH4A1

GeneID: <u>8659</u>

NCBI Accession: NP_003739.2

KEGG: hsa:8659

String: <u>9606.ENSP00000364490</u>

Molecular Weight: Calculated MW: 62 kDa

Observed MW: 62 kDa

Buffer: PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC,

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.