



StAR Related Lipid Transfer Domain Containing 8 (STARD8) Antibody

Catalogue No.: abx238293



WB analysis of HEK-293 cells, using STARD8 antibody (1/400 dilution).



IHC-P analysis of human stomach cancer tissue, using STARD8 Antibody (1/50 dilution).

STARD8 Antibody is a Rabbit Polyclonal against STARD8.

Target: StAR Related Lipid Transfer Domain Containing 8 (STARD8)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB, IHC

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000, IHC: 1/20 - 1/200. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: StAR-related lipid transfer (START) domain containing 8

Isotype: IgG

Form: Liquid

Purity: ≥ 95% (SDS-PAGE)

Datasheet

Version: 2.0.0
Revision date: 18 Mar 2025



Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	Q92502 (UniProt , ExPASy)
Gene Symbol:	STARD8
GeneID:	9754
OMIM:	300689
HGNC:	19161
KEGG:	hsa:9754
Ensembl:	ENSG00000130052
String:	9606.ENSP00000363727
Molecular Weight:	Observed MW: 115-120 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.