



Signal Transducer And Activator of Transcription 3 (STAT3) Antibody


Catalogue No.: abx238298



WB analysis of HeLa cells, using STAT3 antibody (1/300 dilution).



IHC-P analysis of human breast cancer tissue, using STAT3 Antibody (1/50 dilution).



IF analysis of HeLa cells, using STAT3 antibody (1/25 dilution) and Rhodamine-Goat anti-Rabbit IgG.

STAT3 Antibody is a Rabbit Polyclonal against STAT3.

Target: Signal Transducer And Activator of Transcription 3 (STAT3)**Clonality:** Polyclonal**Reactivity:** Human, Mouse, Rat**Tested Applications:** ELISA, WB, IHC, IF/ICC**Host:** Rabbit**Recommended dilutions:** WB: 1/200 - 1/500, IHC: 1/50 - 1/200, IF/ICC: 1/10 - 1/100. Optimal dilutions/concentrations should be determined by the end user.**Conjugation:** Unconjugated

Datasheet

Version: 2.0.0
Revision date: 13 Aug 2025



Immunogen:	signal transducer and activator of transcription 3 (acute-phase response factor)
Isotype:	IgG
Form:	Liquid
Purity:	≥ 95% (SDS-PAGE)
Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	P40763 (UniProt , ExPASy)
KEGG:	hsa:6774
String:	9606.ENSP00000264657
Molecular Weight:	Observed MW: 85 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.