



Mitogen-Activated Protein Kinase Kinase Kinase 7 (MAP3K7) Antibody

Catalogue No.: abx238492



WB analysis of CHO cells, using MAP3K7 antibody (1/1000 dilution).



IHC-P analysis of human prostate cancer tissue, using MAP3K7 Antibody (1/200 dilution). Heat-mediated antigen retrieval was carried out with Tris-EDTA buffer, pH 9.

TAK1 Antibody is a Rabbit Polyclonal against TAK1.

Target: Mitogen-Activated Protein Kinase Kinase Kinase 7 (MAP3K7)

Clonality: Polyclonal

Reactivity: Human, Mouse, Hamster

Tested Applications: ELISA, WB, IHC

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000, IHC: 1/20 - 1/200. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: mitogen-activated protein kinase kinase kinase 7

Isotype: IgG

Form: Liquid

Purity: ≥ 95% (SDS-PAGE)

Datasheet

Version: 3.0.0
Revision date: 27 Aug 2025



Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	P49116 (UniProt , ExPASy)
Gene Symbol:	NR2C2
GeneID:	7182
OMIM:	601426
HGNC:	7972
KEGG:	hsa:7182
Ensembl:	ENSG00000177463
String:	9606.ENSP00000483059
Molecular Weight:	Observed MW: 64-67 kDa, 75-80 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.