



Tubulin Folding Cofactor A (TBCA) Antibody

Catalogue No.: abx238516



WB analysis of human brain tissue, using TBCA antibody (1/500 dilution).



IHC-P analysis of human prostate cancer tissue, using TBCA antibody (1/50 dilution).

TBCA Antibody is a Rabbit Polyclonal against TBCA.

Target: Tubulin Folding Cofactor A (TBCA)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB, IHC

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000, IHC: 1/20 - 1/200. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: tubulin folding cofactor A

Isotype: IgG

Form: Liquid

Purity: ≥ 95% (SDS-PAGE)

Datasheet

Version: 2.0.0
Revision date: 13 Sep 2025



Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	O75347 (UniProt , ExPASy)
Gene Symbol:	TBCA
GeneID:	6902
OMIM:	610058
HGNC:	11579
KEGG:	hsa:6902
Ensembl:	ENSG00000171530
String:	9606.ENSP00000306362
Molecular Weight:	Observed MW: 13 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.