



Voltage-Dependent Anion-Selective Channel Protein 3 (VDAC3) Antibody


Catalogue No.: abx239389



WB analysis of mouse testis tissue, using VDAC3 antibody (1/300 dilution).



IHC-P analysis of human testis tissue, using VDAC3 antibody (1/50 dilution).



IP analysis of mouse testis tissue lysate (4800 µg), using VDAC3 antibody (4 µg, detection: 1/300 dilution).

VDAC3 Antibody is a Rabbit Polyclonal against VDAC3.

Target: Voltage-Dependent Anion-Selective Channel Protein 3 (VDAC3)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB, IHC, IP

Host: Rabbit

Recommended dilutions: WB: 1/200 - 1/2000, IHC: 1/20 - 1/200, IP: 1/200 - 1/1000. Optimal dilutions/concentrations should be determined by the end user.

Datasheet

Version: 2.0.0
Revision date: 27 Aug 2025



Conjugation:	Unconjugated
Immunogen:	voltage-dependent anion channel 3
Isotype:	IgG
Form:	Liquid
Purity:	≥ 95% (SDS-PAGE)
Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	Q9Y277 (UniProt , ExPASy)
Gene Symbol:	VDAC3
GeneID:	7419
OMIM:	610029
HGNC:	12674
Ensembl:	ENSG00000078668
String:	9606.ENSP00000428845
Molecular Weight:	Observed MW: 23-27 kDa, 45 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.