


Vesicle Transport Through Interaction With T-SNAREs Homolog 1A (VTI1A) Antibody

Catalogue No.: abx239462



WB analysis of human brain tissue, using VTI1A antibody (1/500 dilution).

IP analysis of MCF-7 cell lysate (520 µg), using VTI1A antibody (3 µg, detection: 1/1000 dilution).

VTI1A Antibody is a Rabbit Polyclonal against VTI1A.

Target:	Vesicle Transport Through Interaction With T-SNAREs Homolog 1A (VTI1A)
Clonality:	Polyclonal
Reactivity:	Human, Mouse, Rat
Tested Applications:	ELISA, WB, IF/ICC, IP
Host:	Rabbit
Recommended dilutions:	WB: 1/1000 - 1/10000, IF/ICC: 1/10 - 1/100, IP: 1/500 - 1/5000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	vesicle transport through interaction with t-SNAREs homolog 1A (yeast)
Isotype:	IgG
Form:	Liquid

Datasheet

Version: 3.0.0
Revision date: 30 Aug 2025



Purity:	≥ 95% (SDS-PAGE)
Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	Q96AJ9 (UniProt , ExPASy)
Gene Symbol:	VTI1A
GeneID:	143187
OMIM:	614316
HGNC:	17792
KEGG:	hsa:143187
Ensembl:	ENSG00000151532
String:	9606.ENSP00000376792
Molecular Weight:	Observed MW: 47 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.