



Von Willebrand Factor A Domain Containing Protein 2 (VWA2) Antibody


Catalogue No.: abx239465



WB analysis of A375 cells, using VWA2 antibody (1/600 dilution).



IHC-P analysis of human colon tissue, using VWA2 Antibody (1/50 dilution).



IF analysis of A375 cells, using VWA2 antibody (1/50 dilution) and AF488-conjugated Goat Anti-Rabbit IgG (H+L).

VWA2 Antibody is a Rabbit Polyclonal against VWA2.

Target: Von Willebrand Factor A Domain Containing Protein 2 (VWA2)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: ELISA, WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 1/200 - 1/2000, IHC: 1/20 - 1/200, IF/ICC: 1/20 - 1/200. Optimal dilutions/concentrations should be determined by the end user.

Datasheet

Version: 2.0.0
Revision date: 22 Aug 2025



Conjugation:	Unconjugated
Immunogen:	von Willebrand factor A domain containing 2
Isotype:	IgG
Form:	Liquid
Purity:	≥ 95% (SDS-PAGE)
Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	Q5GFL6 (UniProt , ExPASy)
Gene Symbol:	VWA2
GeneID:	340706
HGNC:	24709
KEGG:	hsa:340706
Ensembl:	ENSG00000165816
String:	9606.ENSP00000376708
Molecular Weight:	Observed MW: 82 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.