



Wiskott-Aldrich Syndrome Protein (WAS) Antibody


Catalogue No.: abx239471



WB analysis of Raji cells, using WAS antibody (1/600 dilution).



IHC-P analysis of human spleen tissue, using WAS Antibody (1/50 dilution).



IP analysis of Raji cell lysate (3500 µg), using WASP antibody (3 µg, detection: 1/500 dilution).

WASP Antibody is a Rabbit Polyclonal against WASP.

Target: Wiskott-Aldrich Syndrome Protein (WAS)

Clonality: Polyclonal

Reactivity: Human, Mouse

Tested Applications: ELISA, WB, IHC, IP

Host: Rabbit

Recommended dilutions: WB: 1/200 - 1/2000, IHC: 1/20 - 1/200, IP: 1/200 - 1/2000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Datasheet

Version: 3.0.0
Revision date: 30 Aug 2025



Immunogen:	Wiskott-Aldrich syndrome (eczema-thrombocytopenia)
Isotype:	IgG
Form:	Liquid
Purity:	≥ 95% (SDS-PAGE)
Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	P42768 (UniProt , ExPASy)
KEGG:	hsa:7454
String:	9606.ENSP00000365891
Molecular Weight:	Observed MW: 53 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.