


Transcriptional Activator Myb (c-Myb) Antibody

Catalogue No.: abx432520



abx432520 (0.5 µg/ml) staining of Jurkat lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

c-Myb Antibody is a Goat Polyclonal antibody against c-Myb.

Target:	Transcriptional Activator Myb (c-Myb)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	P-ELISA, WB
Host:	Goat
Recommended dilutions:	P-ELISA: 1/8000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx616255 - Internal region, 557-569 AA: C-HRDQTVKTQKENT
Isotype:	IgG
Form:	Liquid
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Gene Symbol:	MYB
GeneID:	4602
NCBI Accession:	NP_001123645.1, NP_005366.2, NP_001123644.1, NP_001155128.1, NP_001155129.1, NP_001155130.1, NP_001155131.1, NP_001155132.1

Datasheet

Version: 3.0.0

Revision date: 06 Mar 2025



Buffer: Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.

Concentration: 0.5 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only