


2-Acylglycerol O-Acyltransferase 3 (MOGT3) Antibody

Catalogue No.: abx026523



Acyl-CoA:monoacylglycerol acyltransferase (MOGAT; EC 2.3.1.22) catalyzes the synthesis of diacylglycerol from 2-monoacylglycerol and fatty acyl-CoA (Cheng et al., 2003 [PubMed 12618427]).

Target: 2-Acylglycerol O-Acyltransferase 3 (MOGT3)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: ELISA, WB, FCM

Host: Rabbit

Recommended dilutions: WB: 1/1000, FCM: 1/10 - 1/50. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: KLH-conjugated synthetic peptide between 295-329 amino acids from the C-terminal region of human MOGT3.

Isotype: IgG

Form: Liquid

Datasheet

Version: 3.0.0
Revision date: 10 Jun 2025



Purification:	Purified through a protein A column, followed by peptide affinity purification.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q86VF5 (UniProt , ExPASy)
Gene Symbol:	MOGAT3
KEGG:	hsa:346606
String:	9606.ENSP00000223114
Molecular Weight:	Calculated MW: 38.7 kDa
Buffer:	PBS containing 0.09% sodium azide.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only