



Dipeptidyl Peptidase 4 / CD26 (DPP4) Antibody


Catalogue No.: abx430065




abx430065 (0.3 µg/ml) staining of Rat Lung lysate (35 µg protein in RIPA buffer). Detected by chemiluminescence.



abx430065 (8 µg/ml) staining of paraffin embedded Human Prostate. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



Negative Control showing staining of paraffin embedded Human Prostate, with no primary antibody.



Flow cytometry analysis of paraformaldehyde fixed HepG2 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10 µg/ml) followed by AF488 secondary antibody (1 µg/ml). IgG control: Unimmunized goat IgG (black line) followed by AF488 secondary antibody.

DPP4 Antibody is a Goat Polyclonal antibody against DPP4.

Target: Dipeptidyl Peptidase 4 / CD26 (DPP4)

Clonality: Polyclonal

Reactivity: Human, Rat

Datasheet

Version: 2.0.0
Revision date: 04 Nov 2025



Tested Applications: P-ELISA, WB, IHC, FCM

Host: Goat

Recommended dilutions: P-ELISA: 1/64000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: [abx615170](#) - Internal region: C-PPHFDKSKKYP

Isotype: IgG

Form: Liquid

Purification: Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Gene Symbol: DPP4

GeneID: [1803](#) [13482](#) [25253](#)

NCBI Accession: NP_001926.2

Buffer: Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.

Concentration: 0.5 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.