


Frizzled 2 (FZD2) Antibody


Catalogue No.:abx432024




abx432024 (0.03 µg/ml) staining of Human Heart (A) and (0.1 µg/ml) Placenta (B) lysate (35 µg protein in RIPA buffer). Detected by chemiluminescence.



abx432024 (2.5 µg/ml) staining of paraffin embedded Human Thyroid Gland. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



abx432024 (2.5 µg/ml) staining of paraffin embedded Human Liver. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



abx432024 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 µg/ml) followed by AF488 secondary antibody (2 µg/ml), showing nuclear and cytoplasmic staining. Actin filaments were stained with phalloidin (red). Negative control: Unimmunized goat IgG (10 µg/ml) followed by AF488 secondary antibody (2 µg/ml).

Frizzled 2 Antibody is a Goat Polyclonal antibody against Frizzled 2.

Target: Frizzled 2 (FZD2)

Clonality: Polyclonal

Reactivity: Human

Datasheet

Version: 2.0.0
Revision date: 12 Oct 2025



Tested Applications: P-ELISA, WB, IHC, IF/ICC

Host: Goat

Recommended dilutions: P-ELISA: 1/64000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: [abx615441](#) - Internal region: RYATLEHPFHCPR

Isotype: IgG

Form: Liquid

Purification: Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Gene Symbol: FZD2

GeneID: [2535](#) [57265](#) [64512](#)

NCBI Accession: NP_001457.1

Buffer: Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.

Concentration: 0.5 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.