


## JmjC Domain-Containing Histone Demethylation Protein 2B (JHDM2b) Antibody

Catalogue No.:abx026585



Covalent modification of histones plays critical role in regulating chromatin structure and transcription. While most covalent histone modifications are reversible, only recently has it been established that methyl groups are subject to enzymatic removal from histones. A family of novel JmjC domain-containing histone demethylation (JHDM) enzymes have been identified that perform this specific function. Histone demethylation by JHDM proteins requires cofactors Fe (II) and alpha-ketoglutarate. Family members include JHDM1 (demethylating histone 3 at lysine 36), and JHDM2A as well as JMJD2CH3K9 (both of which demethylate histone 3 at lysine 9). Contributions of histone demethylase activity to tumor development, decreases in cell proliferation, and hormone-dependent transcriptional activation have been observed.

Target: JmjC Domain-Containing Histone Demethylation Protein 2B (JHDM2b)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: ELISA, WB, FCM

Host: Rabbit

Recommended dilutions: WB: 1/1000, FCM: 1/10 - 1/50. Optimal dilutions/concentrations should be determined by the end

user.

Conjugation: Unconjugated

Immunogen: KLH-conjugated synthetic peptide between 869-897 amino acids from the Central region of human

JHDM2b.

## **Datasheet**

Version: 4.0.0 Revision date: 05 Mar 2025



Isotype: IgG

Form: Liquid

**Purification:** Purified Rabbit Polyclonal Antibody.

**Storage:** Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q7LBC6 (UniProt, ExPASy)

**KEGG:** hsa:51780

String: <u>9606.ENSP00000326563</u>

Molecular Weight: Calculated MW: 192 kDa

**Buffer:** PBS containing 0.09% sodium azide.

**Specificity:** Predicted to react with Mouse KDM3B.

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC,

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.