



GTRAP3-18 Antibody

Catalogue No.: abx430883




abx430883 staining (0.1 µg/ml) of Human Brain lysate (RIPA buffer, 35 µg total protein per lane). Primary incubated for 1 hour. Detected by chemiluminescence.




abx430883 (1 µg/ml) staining of Human Cerebellum lysate (35 µg protein in RIPA buffer). Detected by chemiluminescence.



abx430883 (1 µg/ml) staining of K562 cell lysate (35 µg protein in RIPA buffer). Detected by chemiluminescence.




abx430883 (3.75 µg/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.




abx430883 (3.75 µg/ml) staining of paraffin embedded Human Small Intestine. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

Datasheet

Version: 2.0.0
Revision date: 23 May 2025



Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 µg/ml) followed by AF488 secondary antibody (2 µg/ml), showing plasma membrane and ER/cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 µg/ml) followed by AF488 secondary antibody (2 µg/ml).



Flow cytometry analysis of paraformaldehyde fixed A431 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10 µg/ml) followed by AF488 secondary antibody (1 µg/ml). IgG control: Unimmunized goat IgG (10 µg/ml) followed by AF488 secondary antibody.

GTRAP3-18 Antibody is a Goat Polyclonal antibody against GTRAP3-18.

| | |
|-------------------------------|---|
| Target: | GTRAP3-18 |
| Clonality: | Polyclonal |
| Reactivity: | Human |
| Tested Applications: | P-ELISA, WB, IHC, IF/ICC, FCM |
| Host: | Goat |
| Recommended dilutions: | P-ELISA: 1/32000. Optimal dilutions/concentrations should be determined by the end user. |
| Conjugation: | Unconjugated |
| Immunogen: | abx269234 - C-Terminus: C-NRLTDYISKVKE |
| Isotype: | IgG |
| Form: | Liquid |
| Purification: | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. |
| Storage: | Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles. |

Datasheet

Version: 2.0.0
Revision date: 23 May 2025



Gene Symbol: ARL6IP5

GeneID: [10550](#)

NCBI Accession: NP_006398.1

Buffer: Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.

Concentration: 0.5 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only