


Hypermethylated In Cancer 2 Protein (HIC2) Antibody

Catalogue No.: abx432033



abx432033 (0.1 µg/ml) staining of HepG2 lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

HIC2 Antibody is a Goat Polyclonal antibody against HIC2.

Target: Hypermethylated In Cancer 2 Protein (HIC2)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: P-ELISA, WB

Host: Goat

Recommended dilutions: P-ELISA: 1/128000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: [abx616882](#) - Internal region, 188-200 AA: C-PQELPQAKGSDDE

Isotype: IgG

Form: Liquid

Purification: Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Gene Symbol: HIC2

GeneID: [23119](#) [58180](#) [287940](#)

NCBI Accession: NP_055909.2

Datasheet

Version: 4.0.0

Revision date: 10 Jun 2025



Buffer: Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.

Concentration: 0.5 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only