


NCAM1 (Isoform 2) Antibody


Catalogue No.: abx430124



Western blot analysis of Human Frontal Cortex (A) and Cerebellum (B) lysate (35 µg protein in RIPA buffer) using NCAM1 antibody ((A): 0.01 µg/ml, (B): 0.1 µg/ml).



IHC-P analysis of paraffin-embedded Human Adrenal Gland using NCAM1 antibody (5 µg/ml).



Flow cytometry analysis of paraformaldehyde fixed U251 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10 µg/ml) followed by AF488 secondary antibody (1 µg/ml). IgG control: Unimmunized goat IgG (black line) followed by AF488 secondary antibody.

NCAM1 (Isoform 2) Antibody is a Goat Polyclonal antibody against NCAM1 (Isoform 2).

Target: NCAM1 (Isoform 2)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: P-ELISA, WB, IHC, FCM

Host: Goat

Recommended dilutions: P-ELISA: 1/32000, WB: 0.03-0.3 µg/ml, IHC: 5 µg/ml, FCM: 10 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Datasheet

Version: 3.0.0
Revision date: 12 Jun 2025



Immunogen:	abx616596 - Internal region: C-KASWTRPEKQETLD
Isotype:	IgG
Form:	Liquid
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Gene Symbol:	NCAM1
GeneID:	4684 24586
NCBI Accession:	NP_851996.2
Buffer:	Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.
Concentration:	0.5 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.