



Neural Cell Adhesion Molecule 2 (NCAM2) Antibody


Catalogue No.: abx430375



24-h analysis of a PFA-perfused cryosection of Mouse Olfactory bulb, using Neural Cell Adhesion Molecule 2 (NCAM2) Antibody (0.5 µg/ml). Antigen retrieval with methanol for 10 mins, followed by 1% SDS and IF analysis.



48-h analysis of a PFA-perfused cryosection of Mouse Olfactory bulb, using Neural Cell Adhesion Molecule 2 (NCAM2) Antibody (0.5 µg/ml). Antigen retrieval with citrate buffer for 10 mins, followed by IF analysis using NCAM2 (red), VGLUT2 (blue) and NQO1 (green).



IHC-P analysis of Human cortex tissue, using Neural Cell Adhesion Molecule 2 (NCAM2) Antibody (3.75 µg/ml).

NCAM2 Antibody is a Goat Polyclonal antibody against NCAM2.

Target: Neural Cell Adhesion Molecule 2 (NCAM2)

Clonality: Polyclonal

Reactivity: Human, Mouse

Tested Applications: P-ELISA, WB, IHC

Host: Goat

Recommended dilutions: P-ELISA: 1/4000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Datasheet

Version: 3.0.0
Revision date: 03 Nov 2025



Immunogen:	abx269790 - C-Terminus: C-KVSNDIIQSKEDDSK
Isotype:	IgG
Form:	Liquid
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Gene Symbol:	NCAM2
GeneID:	4685 17968 288280
NCBI Accession:	NP_004531.2
Buffer:	Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.
Concentration:	0.5 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only