


Nuclear Factor of Activated T-Cells 3 (NFATC3) Antibody

Catalogue No.: abx430730



abx430730 (0.03 µg/ml) staining of Jurkat nuclear lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

NFATC3 Antibody is a Goat Polyclonal antibody against NFATC3.

Target:	Nuclear Factor of Activated T-Cells 3 (NFATC3)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	P-ELISA, WB
Host:	Goat
Recommended dilutions:	P-ELISA: 1/16000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx615976 - Internal region (near N-Terminus): C-HQELDAHEDDLQIN
Isotype:	IgG
Form:	Liquid
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Gene Symbol:	NFATC3
GeneID:	4775
NCBI Accession:	NP_775188.1, NP_004546.1, NP_775186.1, NP_775187.1

Datasheet

Version: 3.0.0

Revision date: 02 Jul 2025



Buffer: Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.

Concentration: 0.5 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only