


Proteasome Activator Complex Subunit 2 (PSME2) Antibody

Catalogue No.: abx431529



abx431529 (0.5 µg/ml) staining of Human Spleen lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



abx431529 (5 µg/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

PSME2 Antibody is a Goat Polyclonal antibody against PSME2.

Target: Proteasome Activator Complex Subunit 2 (PSME2)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: P-ELISA, WB, IHC

Host: Goat

Recommended dilutions: P-ELISA: 1/32000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: [abx615137](#) - C-Terminus: C-NLEKIVNPKGEEKP

Isotype: IgG

Form: Liquid

Purification: Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Datasheet

Version: 4.0.0

Revision date: 01 May 2025



Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q9UL46 (UniProt , ExPASy)
Gene Symbol:	PSME2
GeneID:	5721
OMIM:	602161
NCBI Accession:	NP_002809.2
HGNC:	9569
KEGG:	hsa:5721
Ensembl:	ENSG00000100911
String:	9606.ENSP00000216802
Buffer:	Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.
Concentration:	0.5 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.