


NAD-Dependent Protein Deacetylase Sirtuin-3, Mitochondrial (SIRT3) Antibody

Catalogue No.: abx433275



abx433275 (0.2 µg/ml) staining of Human Liver lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

SIRT3 Antibody is a Goat Polyclonal antibody against SIRT3.

Target:	NAD-Dependent Protein Deacetylase Sirtuin-3, Mitochondrial (SIRT3)
Research Area:	Signal Transduction
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	P-ELISA, WB
Host:	Goat
Recommended dilutions:	P-ELISA: 1/128000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx269781 - C-Terminus: C-QRETGKLDGPKD
Isotype:	IgG
Form:	Liquid
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Gene Symbol:	SIRT3

Datasheet

Version: 4.0.0

Revision date: 09 Sep 2025



GeneID: [23410](#)

NCBI Accession: NP_036371.1, NP_001017524.1

Buffer: Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.

Concentration: 0.5 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only