



Sorting Nexin 18 (SNX18) Antibody


Catalogue No.: abx430604




Western blot analysis of A549 lysate (35 µg protein in RIPA buffer) using Sorting Nexin 18 (SNX18) Antibody (1 µg/ml, 1 hour). Detected by chemiluminescence.



Immunofluorescence analysis of paraformaldehyde fixed HeLa cells permeabilized with 0.15% Triton, using Sorting Nexin 18 (SNX18) Antibody (10 µg/ml, 1 hour) followed by AF488 secondary antibody (2 µg/ml), and DAPI (blue) nuclear stain. Negative control: Unimmunized goat IgG (10 µg/ml) followed by AF488 secondary antibody (2 µg/ml).



Immunofluorescence analysis of paraformaldehyde fixed U2OS cells permeabilized with 0.15% Triton, using Sorting Nexin 18 (SNX18) Antibody (10 µg/ml, 1 hour) followed by AF488 secondary antibody (2 µg/ml), and DAPI (blue) nuclear stain. Negative control: Unimmunized goat IgG (10 µg/ml) followed by AF488 secondary antibody (2 µg/ml).



Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line) permeabilized with 0.5% Triton, using Sorting Nexin 18 (SNX18) Antibody (10 µg/ml, 1 hour) followed by AF488 secondary antibody (1 µg/ml). IgG control: Unimmunized goat IgG (black line) followed by AF488 secondary antibody.

SNX18 Antibody is a Goat Polyclonal antibody against SNX18.

Target: Sorting Nexin 18 (SNX18)

Clonality: Polyclonal

Reactivity: Human

Datasheet

Version: 3.0.0
Revision date: 06 Mar 2025



Tested Applications: P-ELISA, IF/ICC, FCM

Host: Goat

Recommended dilutions: P-ELISA: 1/32000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: [abx269329](#) - C-Terminus: C-KVQRVPLMTVLSF

Isotype: IgG

Form: Liquid

Purification: Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Gene Symbol: SNX18

GeneID: [112574](#)

NCBI Accession: NP_443102.2

Buffer: Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.

Concentration: 0.5 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.