


Signal Transducing Adaptor Molecule 2 (STAM2) Antibody


Catalogue No.: abx432003



Western blot analysis of Human Spleen lysate (35 µg protein in RIPA buffer) using Signal Transducing Adaptor Molecule 2 (STAM2) Antibody (1 µg/ml, 1 hour).



Flow cytometry analysis of paraformaldehyde fixed K562 cells (blue line), permeabilized with 0.5% Triton using Signal Transducing Adaptor Molecule 2 (STAM2) Antibody (10 µg/ml, 1 hour) followed by AF488 secondary antibody (1 µg/ml). IgG control: Unimmunized goat IgG (black line) followed by AF488 secondary antibody.



Flow cytometry analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton using Signal Transducing Adaptor Molecule 2 (STAM2) Antibody (10 µg/ml, 1 hour) followed by AF488 secondary antibody (1 µg/ml). IgG control: Unimmunized goat IgG (black line) followed by AF488 secondary antibody.

STAM2 Antibody is a Goat Polyclonal antibody against STAM2.

Target: Signal Transducing Adaptor Molecule 2 (STAM2)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: P-ELISA, WB, FCM

Host: Goat

Recommended dilutions: P-ELISA: 1/8000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Datasheet

Version: 3.0.0
Revision date: 13 Aug 2025



Immunogen:	abx269886 - Internal region: C-QSIDPTDSKPDSQD
Isotype:	IgG
Form:	Liquid
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Gene Symbol:	STAM2
GeneID:	10254 56324
NCBI Accession:	NP_005834.3
Buffer:	Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.
Concentration:	0.5 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.