



Signal Transducer And Activator of Transcription 5A (STAT5A) Antibody

Catalogue No.: abx431618



abx431618 (0.1 µg/ml) staining of K562 lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



abx431618 (5 µg/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

STAT5A Antibody is a Goat Polyclonal antibody against STAT5A.

Target: Signal Transducer And Activator of Transcription 5A (STAT5A)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: P-ELISA, WB, IHC

Host: Goat

Recommended dilutions: P-ELISA: 1/8000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: [abx616760](#) - Internal region, 681-692 AA: C-KYYTPVLAKAVD

Isotype: IgG

Form: Liquid

Datasheet

Version: 3.0.0

Revision date: 13 Aug 2025



Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Gene Symbol:	STAT5A
GeneID:	6776 20850 24918
NCBI Accession:	NP_003143.2
Buffer:	Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.
Concentration:	0.5 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only