



Triggering Receptor Expressed On Myeloid Cells 2 (Trem2) Antibody

Catalogue No.: abx432230



Immunofluorescence analysis of paraformaldehyde fixed 3T3-L1 cells permeabilized with 0.15% Triton using Triggering Receptor Expressed On Myeloid Cells 2 (Trem2) Antibody (1 hour, 10 µg/ml) followed by AF488 secondary antibody (2 µg/ml), and DAPI (blue) nuclear stain. Negative control: Unimmunized goat IgG (10 µg/ml) followed by AF488 secondary antibody (2 µg/ml).



Immunofluorescence analysis of paraformaldehyde fixed RAW264.7 cells permeabilized with 0.15% Triton using Triggering Receptor Expressed On Myeloid Cells 2 (Trem2) Antibody (1 hour, 10 µg/ml) followed by AF488 secondary antibody (2 µg/ml), and DAPI (blue) nuclear stain. Negative control: Unimmunized goat IgG (10 µg/ml) followed by AF488 secondary antibody (2 µg/ml).

Trem2 Antibody is a Goat Polyclonal antibody against Trem2.

Target: Triggering Receptor Expressed On Myeloid Cells 2 (Trem2)

Clonality: Polyclonal

Reactivity: Mouse

Tested Applications: P-ELISA, IF/ICC

Host: Goat

Recommended dilutions: P-ELISA: 1/32000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: [abx615208](#) - Internal region: C-QVEHSTSRNQET

Isotype: IgG

Form: Liquid

Purification: Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Datasheet

Version: 2.0.0

Revision date: 19 Sep 2025



Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Gene Symbol:	TREM2
GeneID:	83433 301227
NCBI Accession:	NP_112544.1
Buffer:	Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.
Concentration:	0.5 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only