



Ubiquitin Conjugating Enzyme E2C (UBE2C) Antibody


Catalogue No.:abx430628




abx430628 (2 μ g/ml) staining of HEK293 (A) and HeLa (B) cell lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



abx430628 Immunofluorescence analysis of paraformal dehyde fixed MCF7 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 µg/ml) followed by AF488 secondary antibody (4 µg/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 µg/ml) followed by AF488 secondary antibody (4 µg/ml).



abx430628 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 μ g/ml) followed by AF488 secondary antibody (4 μ g/ml), showing cytoplasmic /Plasma Membrane staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 μ g/ml) followed by AF488 secondary antibody (4 μ g/ml).



Flow cytometry analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10 µg/ml) followed by AF488 secondary antibody (2 µg/ml). IgG control: Unimmunized goat IgG (black line) followed by AF488 secondary antibody.

UBE2C Antibody is a Goat Polyclonal antibody against UBE2C.

Target: Ubiquitin Conjugating Enzyme E2C (UBE2C)

Clonality: Polyclonal

Reactivity: Human

Datasheet

Version: 2.0.0 Revision date: 28 Aug 2025



Tested Applications: P-ELISA, WB, IF/ICC, FCM, IP

Host: Goat

Recommended dilutions: P-ELISA: 1/32000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: abx269212 - C-Terminus: C-QETYSKQVTSQEP

Isotype: IgG

Form: Liquid

Purification: Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

chromatography using the immunizing peptide.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Gene Symbol: UBE2C

GeneID: <u>11065</u>

NCBI Accession: NP_008950.1, NP_861515.1, NP_861516.1, NP_861517.1, NP_861518.1, NP_001268670.1

Buffer: Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.

Concentration: 0.5 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC,

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.