


Uromodulin (UMOD) Antibody

Catalogue No.: abx433430



Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 µg/ml) followed by AF488 secondary antibody (2 µg/ml), showing membrane and Golgi staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 µg/ml) followed by AF488 secondary antibody (2 µg/ml).

Uromodulin Antibody is a Goat Polyclonal antibody against Uromodulin.

Target:	Uromodulin (UMOD)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	P-ELISA, IF/ICC
Host:	Goat
Recommended dilutions:	P-ELISA: 1/64000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx616868 - Internal region, 525-536 AA: DRCPHTRDSTIQ
Isotype:	IgG
Form:	Liquid
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Gene Symbol:	UMOD
GeneID:	7369
NCBI Accession:	NP_003352.2

Datasheet

Version: 3.0.0

Revision date: 31 Aug 2025



Buffer: Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.

Concentration: 0.5 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only