


Vav 3 Oncogene (VAV3) Antibody


Catalogue No.: abx433439




abx433439 (0.5 µg/ml) staining of Human Bone Marrow lysate (35 µg protein in RIPA buffer). Detected by chemiluminescence.



abx433439 (7.5 µg/ml) staining of paraffin embedded Human Spleen. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



abx433439 Negative Control showing staining of paraffin embedded Human Spleen, with no primary antibody.



Flow cytometry analysis of paraformaldehyde fixed K562 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10 µg/ml) followed by AF488 secondary antibody (1 µg/ml). IgG control: Unimmunized goat IgG (black line) followed by AF488 secondary antibody.

VAV3 Antibody is a Goat Polyclonal antibody against VAV3.

Target: Vav 3 Oncogene (VAV3)

Clonality: Polyclonal

Reactivity: Human

Datasheet

Version: 3.0.0
Revision date: 04 Sep 2025



Tested Applications: P-ELISA, IHC, FCM

Host: Goat

Recommended dilutions: P-ELISA: 1/32000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: [abx269563](#) - Internal region: C-SGEQGTLKLPEK

Isotype: IgG

Form: Liquid

Purification: Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Gene Symbol: VAV3

GeneID: [10451](#)

NCBI Accession: NP_006104.4, NP_001073343.1

Buffer: Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.

Concentration: 0.5 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.