



Voltage-Dependent Anion-Selective Channel Protein 2 (VDAC2) Antibody


Catalogue No.: abx433441



Western blot analysis of Human Cerebellum lysate (35 µg in RIPA buffer) using Voltage-Dependent Anion-Selective Channel Protein 2 (VDAC2) Antibody (0.3 µg/ml).



Western blot analysis of CaCo-2 (A) and LNCaP (B) cell lysate (35 µg in RIPA buffer) using Voltage-Dependent Anion-Selective Channel Protein 2 (VDAC2) Antibody (1 µg/ml).



Immunohistochemistry analysis of paraffin-embedded Rat Liver using Voltage-Dependent Anion-Selective Channel Protein 2 (VDAC2) Antibody (6 µg/ml). Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



Immunohistochemistry analysis of paraffin-embedded Rat Liver (negative control), with no primary antibody.

VDAC2 Antibody is a Goat Polyclonal antibody against VDAC2.

Target: Voltage-Dependent Anion-Selective Channel Protein 2 (VDAC2)

Clonality: Polyclonal

Datasheet

Version: 2.0.0
Revision date: 02 Sep 2025



Reactivity:	Human
Tested Applications:	P-ELISA, WB, IHC
Host:	Goat
Recommended dilutions:	P-ELISA: 1/2000, WB: 0.3-1 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx269505 - C-Terminus: C-GHKVGLALELEA
Isotype:	IgG
Form:	Liquid
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Gene Symbol:	VDAC2
GeneID:	7417
NCBI Accession:	NP_003366.2
Buffer:	Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.
Concentration:	0.5 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.