


Vesicular Glutamate Transporter 2 (SLC17A6) Antibody

Catalogue No.: abx431246



abx431246 (2 µg/ml) staining of Human Cerebellum lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Vesicular Glutamate Transporter 2 (SLC17A6) Antibody is a Goat Polyclonal antibody against Vesicular Glutamate Transporter 2 (SLC17A6).

Target: Vesicular Glutamate Transporter 2 (SLC17A6)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: P-ELISA, WB

Host: Goat

Recommended dilutions: P-ELISA: 1/2000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: [abx617072](#) - Internal region, 521-534 AA: C-EDELDEETGDITQN

Isotype: IgG

Form: Liquid

Purification: Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Gene Symbol: SLC17A6

GeneID: [57084](#) [140919](#) [84487](#)

Datasheet

Version: 4.0.0

Revision date: 06 Oct 2025



NCBI Accession: NP_065079.1

Buffer: Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.

Concentration: 0.5 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only