



WNT1 Inducible Signaling Pathway Protein 1 / WISP1 (CCN4) Antibody


Catalogue No.: abx431710




HEK293 overexpressing WISP1 with C-terminal tag (DYKDDDDK) and probed with anti-DYKDDDDK in the left panel and with abx431710 in the right panel (mock transfection in first lane in each panel).




abx431710 (0.1 µg/ml) staining of Human Tonsil lysate (35 µg protein in RIPA buffer). Detected by chemiluminescence.



abx431710 (0.3 µg/ml) staining of Caco-2 cell lysate 1 (A) and Caco-2 cell lysate 2 (B) + petide (C). (35 µg protein in RIPA buffer). Detected by chemiluminescence.




Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 µg/ml) followed by AF488 secondary antibody (2 µg/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 µg/ml) followed by AF488 secondary antibody (2 µg/ml).



Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 µg/ml) followed by AF488 secondary antibody (2 µg/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 µg/ml) followed by AF488 secondary antibody (2 µg/ml).

Datasheet

Version: 3.0.0
Revision date: 22 Dec 2025



Flow cytometry analysis of paraformaldehyde fixed A431 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10 µg/ml) followed by AF488 secondary antibody (1 µg/ml). IgG control: Unimmunized goat IgG (black line) followed by AF488 secondary antibody.

WISP1 Antibody is a Goat Polyclonal antibody against WISP1.

Target:	WNT1 Inducible Signaling Pathway Protein 1 / WISP1 (CCN4)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	P-ELISA, WB, IF/ICC, FCM
Host:	Goat
Recommended dilutions:	P-ELISA: 1/128000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx615336 - C-Terminus: C-ESYPDFSEIAN
Isotype:	IgG
Form:	Liquid
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Gene Symbol:	CCN4
GeneID:	8840 22402 65154
NCBI Accession:	NP_003873.1, NP_543028.1
Buffer:	Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.

Datasheet

Version: 3.0.0

Revision date: 22 Dec 2025



Concentration: 0.5 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only