


RAC-Alpha Serine/threonine-Protein Kinase Phospho-Ser473 (AKT1 pS473) Antibody

Catalogue No.: abx000166



Western blot analysis of extracts of Jurkat cells, using Phospho-AKT1-S473 antibody (abx000166) at 1/2000 dilution. Jurkat cells were treated by Calyculin A (100nM) for 30 minutes.

AKT1 (pS473) Antibody is a Rabbit Polyclonal antibody against AKT1 (pS473). The serine-threonine protein kinase encoded by the AKT1 gene is catalytically inactive in serum-starved primary and immortalized fibroblasts. AKT1 and the related AKT2 are activated by platelet-derived growth factor. The activation is rapid and specific, and it is abrogated by mutations in the pleckstrin homology domain of AKT1. It was shown that the activation occurs through phosphatidylinositol 3-kinase. In the developing nervous system AKT is a critical mediator of growth factor-induced neuronal survival. Survival factors can suppress apoptosis in a transcription-independent manner by activating the serine/threonine kinase AKT1, which then phosphorylates and inactivates components of the apoptotic machinery. Mutations in this gene have been associated with the Proteus syndrome. Multiple alternatively spliced transcript variants have been found for this gene.

Target: RAC-Alpha Serine/threonine-Protein Kinase Phospho-Ser473 (AKT1 pS473)

Clonality: Polyclonal

Target Modification: Ser473

Modification: Phosphorylation

Reactivity: Human, Mouse, Rat

Tested Applications: WB

Host: Rabbit

Recommended dilutions: WB: 1/100 - 1/500. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: A synthetic phosphorylated peptide around S473 of human Akt1.

Isotype: IgG

Form: Liquid

Datasheet

Version: 2.0.0

Revision date: 23 May 2024



Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P31749 (UniProt , ExPASy)
Gene Symbol:	AKT1
GeneID:	207
NCBI Accession:	NP_005154.2
KEGG:	hsa:207
String:	9606.ENSP00000451828
Molecular Weight:	Calculated MW: 56 kDa Observed MW: 60 kDa
Buffer:	PBS, pH 7.3, containing 0.05% Proclin-300, 50% glycerol.
Concentration:	> 0.2 mg/ml
Note:	This product is for research use only.

For Reference Only