



Cancer/Testis Antigen 1A (CTAG1A) Antibody


Catalogue No.: abx027441



WB analysis of HT-1080 whole cell lysates.



WB analysis of HT-1080 and U266B1 whole cell lysates.



IHC-P analysis of human gastric carcinoma tissue, using CTAG1A antibody (1/250 dilution).

CTAG1A is an identical copy of the CTAG1B gene (MIM 300156) (Aradhya et al., 2001 [PubMed 11709543]).[supplied by OMIM].

Target: Cancer/Testis Antigen 1A (CTAG1A)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: ELISA, WB, IHC

Host: Rabbit

Recommended dilutions: WB: 1/1000, IHC-P: 1/250. Not tested in IHC-F. Optimal dilutions/concentrations should be determined by the end user.

Datasheet

Version: 3.0.0
Revision date: 09 Jun 2025



Conjugation:	Unconjugated
Immunogen:	KLH-conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human CTAG1A.
Isotype:	IgG
Form:	Liquid
Purification:	Purified through a protein A column, followed by peptide affinity purification.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P78358 (UniProt , ExPASy)
Gene Symbol:	CTAG1A
GeneID:	1485
KEGG:	hsa:1485, hsa:246100
String:	9606.ENSP00000469441
Molecular Weight:	Calculated MW: 18 kDa
Buffer:	PBS containing 0.09% sodium azide.
Concentration:	0.45 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.